

# Physiologic basis for understanding quantitative dehydration assessment<sup>1–4</sup>

Samuel N Cheuvront, Robert W Kenefick, Nisha Charkoudian, and Michael N Sawka

## ABSTRACT

Dehydration (body water deficit) is a physiologic state that can have profound implications for human health and performance. Unfortunately, dehydration can be difficult to assess, and there is no single, universal gold standard for decision making. In this article, we review the physiologic basis for understanding quantitative dehydration assessment. We highlight how phenomenologic interpretations of dehydration depend critically on the type (dehydration compared with volume depletion) and magnitude (moderate compared with severe) of dehydration, which in turn influence the osmotic (plasma osmolality) and blood volume–dependent compensatory thresholds for antidiuretic and thirst responses. In particular, we review new findings regarding the biological variation in osmotic responses to dehydration and discuss how this variation can help provide a quantitative and clinically relevant link between the physiology and phenomenology of dehydration. Practical measures with empirical thresholds are provided as a starting point for improving the practice of dehydration assessment. *Am J Clin Nutr* 2013;97:455–62.

## INTRODUCTION

Dehydration (body water deficit) is a common physiologic state that can have profound implications for human health (1–7) and performance (8). Although mild dehydration can be easily corrected and is principally associated with impaired physical performance (8), it may be linked with common public health disorders if left chronically untreated (9, 10). A greater severity of dehydration can result in significant medical costs, morbidity, and mortality across the life span (11, 12). Although the physiology of osmotic and vascular volume responses to dehydration in humans have been well described (13, 14), the phenomenology of dehydration assessment has not. For example, there is no single, universal gold standard method of dehydration assessment for clinical decision making (7, 15, 16), which contributes greatly to the difficulty that clinicians encounter when trying to accurately assess dehydration in practice (17–25). This discordance between the physiology and phenomenology of dehydration is a recognized source of clinical confusion (17) for which clarity is needed to improve the practice of dehydration assessment.

In this review, we highlight how phenomenologic interpretations of dehydration depend critically on the type (dehydration

compared with volume depletion) and magnitude (moderate compared with severe) of dehydration, which, in turn, influence the plasma osmolality (Posm)<sup>5</sup>– and blood volume (BV)–dependent compensatory thresholds for antidiuretic and thirst responses. We also discuss the recent application of biological variation analysis to osmotic responses during dehydration for its novel potential as an adjunct (17) to clinical decision making. Posm is the primary focus of this review because it is the key regulated variable in fluid balance (13, 14, 26–28), and it is commonly used to screen for dehydration and complement more quantitative differential diagnoses of dysnatremias and other diseases (3, 5, 28–30). The osmolality of other body fluids commonly used to assess dehydration (ie, urine and saliva) are also mentioned as is the practical assessment of volume depletion. Descriptions of other potential methods of dehydration and volume-depletion assessment have been provided by other authors (7, 16, 19, 31, 32). Complementary reviews (33) are similarly suggested for detailed information related to sodium (natriuresis and appetite) and nonosmotic contributors (eg, baroreceptors) to osmotic homeostasis.

## FUNDAMENTALS OF OSMOTIC RESPONSES TO DEHYDRATION IN HUMANS

In its simplest form, the net body water balance is generally the zero sum of food (water and solute) and fluid intake minus insensible and obligatory renal water losses (7). Fluid intakes,

<sup>1</sup> From the US Army Research Institute of Environmental Medicine, Natick, MA.

<sup>2</sup> The opinions or assertions contained herein are the private views of the authors and should not be construed as official or reflecting the views of the Army or the Department of Defense.

<sup>3</sup> Supported by the United States Army Medical Research and Materiel Command.

<sup>4</sup> Address correspondence to SN Cheuvront, Thermal and Mountain Medicine Division, US Army Research Institute of Environmental Medicine, Kansas Street, Building 42, Natick, MA 01760-5007. E-mail: samuel.n.cheuvront@us.army.mil.

<sup>5</sup> Abbreviations used: AVP, arginine vasopressin; Bm, body mass; BV, blood volume; Posm, plasma osmolality; PV, plasma volume; Sosm, saliva osmolality; TBW, total body water; Uosm, urine osmolality.

Received June 5, 2012. Accepted for publication December 14, 2012.

First published online January 23, 2013; doi: 10.3945/ajcn.112.044172.

Report Documentation Page			Form Approved OMB No. 0704-0188		
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE <b>2012</b>		2. REPORT TYPE		3. DATES COVERED	
4. TITLE AND SUBTITLE <b>Physiologic Basis for Understanding Quantitative Dehydration Assessment</b>		5a. CONTRACT NUMBER			
		5b. GRANT NUMBER			
		5c. PROGRAM ELEMENT NUMBER			
6. AUTHOR(S) <b>Samuel Cheuvront; Robert Kenefick; Nisha Charkoudian; Michael Sawka</b>		5d. PROJECT NUMBER			
		5e. TASK NUMBER			
		5f. WORK UNIT NUMBER			
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>Thermal and Mountain Medicine Division,U.S. Army Research Institute of Environmental Medicine,Natick,MA,01760-5007</b>		8. PERFORMING ORGANIZATION REPORT NUMBER <b>MISC 13-22</b>			
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)		10. SPONSOR/MONITOR'S ACRONYM(S)			
		11. SPONSOR/MONITOR'S REPORT NUMBER(S)			
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release; distribution unlimited.</b>					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT <b>Dehydration (body water deficit) is a physiologic state that can have profound implications for human health and performance. Unfortunately, dehydration can be difficult to assess, and there is no single, universal gold standard for decision making. In this article, we review the physiologic basis for understanding quantitative dehydration assessment. We highlight how phenomenologic interpretations of dehydration depend critically on the type (dehydration compared with volume depletion) and magnitude (moderate compared with severe) of dehydration, which in turn influence the osmotic (plasma osmolality) and blood volume?dependent compensatory thresholds for antidiuretic and thirst responses. In particular, we review new findings regarding the biological variation in osmotic responses to dehydration and discuss how this variation can help provide a quantitative and clinically relevant link between the physiology and phenomenology of dehydration. Practical measures with empirical thresholds are provided as a starting point for improving the practice of dehydration assessment.</b>					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES <b>8</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			

losses, and needs vary widely in free-living people and are governed heavily by physical activity, environmental stress, and cultural and habitual cues (7, 8, 34–36). Under conditions of ordinary daily body water flux, osmotic constancy is maintained by the secretion of the antidiuretic hormone arginine vasopressin (AVP), which directly influences renal water excretion and conservation in response to intravascular fluid shifts (that result from thermal and positional changes) and ad libitum food and fluid intakes (14, 26, 37–39). Thus, Posm remains stable as the kidneys modify urine osmolal and water excretion in accordance with ordinary living conditions. When a body water deficit in excess of ordinary flux occurs (dehydration), threshold increases in Posm (primary) and decreases in BV (secondary) produce compensatory water-conservation (renal) and water-acquisition (thirst) responses (14, 26). As a result, the discriminatory power of renal excretion measures for the detection of dehydration is always secondary to changes in Posm (28).

AVP is synthesized in supraoptic and paraventricular nuclei of the hypothalamus and is released from the posterior pituitary (14, 26). Basal AVP concentrations can fluctuate considerably in response to ordinary postural and skin-temperature (skin blood flow) shifts in BV (39). However, a threshold reduction in BV >10% is required to elicit greater (compensatory) AVP secretion, whereas smaller reductions in BV primarily act to enhance the sensitivity of the AVP response to changes in Posm (40–42). Osmotic homeostasis (<1–2% deviation in Posm) is also maintained by basal AVP regulation, but compared to BV smaller threshold increases in Posm (>2%) produce intracellular dehydration and compensatory increases in AVP secretion, renal water conservation, and thirst (14, 43).

When the net balance between water intake and output becomes negative (dehydration), renal water conservation is insufficient to restore fluid balance. Obligatory renal water losses persist, and fluid acquisition must occur, to restore the body water balance (28, 44). However, the Posm threshold for thirst is highly variable in people (27, 45, 46), and thirst mechanisms are subject to numerous influences unrelated to the body water balance (47). In humans, fluid losses (because of sweating, vomiting, or diarrhea) can easily outpace oral intakes. Peripheral osmoreceptors (eg, gut) (14) and oropharyngeal cues trigger thirst satiety well before volume is fully restored (26, 48, 49), even when dehydration is substantial (50). This transitory response acts to buffer the presystemic impact of ingested fluids (14) but often leads to involuntary dehydration when water is consumed without food (solute) (47, 51).

## TWO CRITICAL CAVEATS TO UNDERSTANDING OSMOTIC RESPONSES TO DEHYDRATION

### Caveat 1: a sufficient body water–deficit threshold must be reached before compensatory reactions become reliably engaged

Percentage reductions in body mass (Bm) that exceed typical human variation are depicted in **Figure 1**, whereby a change in Bm is equated with a change in total body water (TBW). The change in Bm is used as the criterion value for practical purposes but also because the random measurement error for tracer-dilution methods (the change in TBW) is larger than the same for Bm (52). Typical human variation is defined as the day-to-day CV in Bm, which is <1.0% when fluid intake and activity

are tightly controlled (53, 54). As a consequence, day-to-day change in Bm must exceed 1% and approach 2% (ie,  $\sqrt{2} \times 1.65$ ) to be considered truly atypical ( $P < 0.05$ ; 1-tailed test). Therefore, day-to-day fluctuations in Bm <1–2% cannot be reliably associated with perturbations in body water beyond ordinary (sinusoidal) physiologic and behavioral body water regulation (55). Under these circumstances, renal water excretion or conservation is a reflection of the flux produced by fluctuating AVP concentrations in response to widely ranging dietary fluid intakes, osmolar loads, and ordinary compartmental fluid shifts without discernible changes in TBW, Posm, or, by extension, intracellular hydration (26, 37–39, 56). Thus, day-to-day fluctuations in Bm or TBW within this range should be interpreted as euhydrated (the state of normal hydration).


### Caveat 2: body water–deficit threshold for dehydration depends critically on the type and magnitude of the body water deficit incurred

A 2% increase in Posm ( $\sim 5$  mmol/kg) and a 10% decrease in BV ( $\sim 0.5$  L) are commonly quoted physiologic thresholds for compensatory water conservation and acquisition (**Figure 1**) (26, 40, 43). Posm increases to greater than  $\sim 5$  mmol/kg in response to dehydration via sweat losses, fluid restriction, or osmotic diarrhea (hypertonic hypovolemia) when those losses exceed  $\sim 2\%$  of Bm (1.4 L at 70 kg) (18, 54, 57–59), which is a threshold that is also consistent with negative physiologic outcomes (7, 8, 60). The variation in sweat sodium losses in people may (61) or may not (62) add uncertainty to the magnitude of the osmotic response to a given water deficit, depending on the delicate balance between sweating rate and sweat sodium concentrations, whereby

$$\text{Volume} \times \text{concentration} = \text{content} \quad (1)$$

Similar considerations may be made of alterations in extracellular volume [plasma volume (PV)], but on the basis of the regression equations shown in **Figure 2**, the anticipated decrease in PV is only  $\sim 0.14$  L at  $\sim 2\%$  dehydration (63) because of the rapid osmotic redistribution of water from the intracellular to the extracellular (interstitial and intravascular) fluid compartment (61, 64). Therefore, hypertonic hypovolemia results in a small ratio of plasma-to-TBW losses ( $\sim 1:10$ ). Hypertonic hypovolemia would not produce intravascular volume losses >10% of BV until a 7% loss of Bm was achieved (**Figure 2**) (63, 65). This effect illustrates the primary influence of Posm as the stimulus for early compensatory water-conservation and -acquisition responses (40–42). Although the osmolalities of other body fluids (eg, urine and saliva) also increase in parallel with Posm and afford a good diagnostic accuracy for dehydration under ideal circumstances (54), they remain secondary (25) and are inferior to Posm for the detection of dehydration for additional reasons.

Isotonic hypovolemia can occur in response to diuretic use, cold or altitude exposure, secretory diarrhea, and vomiting (6, 18, 42, 44, 63, 65–68). The ratio of PV-to-TBW loss is approximately twice as large ( $\sim 1:5$ ) with isotonic hypovolemia than with hypertonic hypovolemia (63, 65, 68). This type of body water loss is often referred to as salt-depletion dehydration (44)



**FIGURE 1.** Body water regulation in response to dehydration. Schematic includes the 2 major types of dehydration, their typical causes, and the estimated magnitude of dehydration required to stimulate a primary osmotic- or volume-dependent response for compensatory water conservation and acquisition (26). A change in TBW was equated with a change in body mass (1 L = 1 kg), whereby dehydration was expressed as a percentage of body mass in accordance with  $(\Delta \text{ body mass} \div \text{body mass}) \times 100$ . Plasma volume and TBW losses are depicted to scale as are their 1:10 and 1:5 ratios for hypertonic and isotonic hypovolemia, respectively. Dashed arrows represent negative feedback. Ang II, angiotensin II; AVP, arginine vasopressin; BV, blood volume; Posm, plasma osmolality; TBW, total body water.

or volume depletion (24, 30) because the added solute loss produces little change in Posm but proportionally greater PV reductions. When there are large losses of solute from the extracellular space, there is a minimal or no osmotic gradient to pull fluids from the larger intracellular space (61, 64). As a result, a smaller ~4% loss of Bm (2.8 L at 70 kg; 0.56-L PV loss) must be incurred to achieve the 10% BV threshold for compensatory water conservation and acquisition with isotonic hypovolemia (Figures 1 and 2).

Methods for the assessment of volume depletion vary widely, with no single, standard approach advocated in the medical or related literature (16, 19). The use of a simple 20-beats/min sit-to-stand heart-rate response provides high specificity but low sensitivity and only marginal diagnostic accuracy even when dehydration is severe (65, 69). BV losses >10% (~1.0 L), whether measured directly (70) or by using lower body negative pressure to simulate equivalent blood losses (71), are required for an improved test sensitivity. The type of water loss with a gastrointestinal illness can be unpredictable or mixed (18, 44), and thus, the presence of both types of dehydration probably


explains much of the difficulty associated with their interpretation (30, 44), in which neither Posm nor BV thresholds for compensatory responses are reached (Figure 1). Under these circumstances, a more heuristic assessment approach is needed (19, 22, 24, 28).

## BIOLOGICAL AND METHODOLOGIC VARIATION

Human variation in osmotic responses to dehydration is primarily biological, but the methodology used to study osmotic responses can also contribute to variation. An understanding and appreciation for these sources of variation can inform probabilistic decision making related to the diagnosis of dehydration (54, 58, 72) and, likely, volume depletion as well (65).

### Threshold and slope of AVP and thirst responses (biology)

It is common to refer to both the threshold and slope of the Posm-AVP relation. The threshold Posm value is associated with the initial increase in AVP secretion above baseline, whereas the slope is the responsiveness (or sensitivity) of the AVP system for



**FIGURE 2.** Linear regression of plasma volume change ( $y$ ) and dehydration [ $\% \Delta$  body mass ( $x$ )] after induction of hypotonic (sweat) or isotonic (diuretic) body water losses. Equations for plasma volume contraction are as follows: diuretic,  $y = -3.8 + 3.6x$ ; sweat,  $y = 1.35 + 2.8x$ . From Cheuvront et al (63).

any given increase in Posm above the threshold value. The osmotic control of AVP, when defined by using slope and sensitivity terms, is highly variable. For example, there appears to be a polygenetic basis for the variation in the slope of the AVP-Posm relation and Posm thresholds for AVP and thirst (46). The relations are highly correlated between monozygotic, but not dizygotic, twin pairs (46). For a healthy and heterogeneous population, the individual AVP-Posm slopes vary 10-fold in individuals but are highly correlated within a subject ( $r = 0.94$ ). The osmotic threshold for AVP varies less in individuals ( $\sim 8$  mmol/kg) but shows only a moderate correlation within subjects ( $r = 0.61$ ).

The individual variation of Posm set points and thresholds for both AVP release and unequivocal thirst relative to what has been commonly reported in the literature for group means is illustrated in **Figure 3** (27). The variation contains the potential influences of sex (73), but not age (74), on osmotic responses. Posm thresholds for AVP release and unequivocal thirst differ in subjects by  $\sim 10$  mmol/kg. The largest difference between Posm thresholds for AVP release and unequivocal thirst within an individual was 17 mmol/kg (subject 5). Also of importance is the difference in the Posm set point relative to the Posm threshold for AVP release and unequivocal thirst; for example, subjects 7 and 15 fell on opposite extremes. Taken together, the data in Figure 3 show that plasma osmotic responses (AVP and thirst) vary considerably in people and have a strong genetic component. These data may partly explain the 20-mmol/kg range in Posm often reported for population reference intervals (eg, 280–300 mmol/kg). Differences in health and hydration states must also contribute to this range, but the volume of fluid ingested and its proximity to measurement can also make an important methodologic contribution (43, 75), even in well-controlled laboratory situations.

### Threshold and slope of AVP and thirst responses (methodology)

Some of the variation in the Posm threshold for AVP and thirst is methodologic rather than physiologic. In this context, moderate water loading is one methodology used to standardize Posm and suppress AVP secretion before imposing an intervention such as saline infusion or water restriction (dehydration). However, this

approach produces low basal Posm values and results in threshold and slope calculations dissimilar from studies in which ad libitum water consumption was permitted before testing (13, 43, 45, 76). Suppressed Posm thresholds for AVP release and thirst in studies that used water-loading methodologies, although experimentally sound, may be unrealistic for free-living people.


Interpreting osmotic responses after water loading should be approached with caution. For example, the application of regression equations for AVP-Posm and AVP-urine osmolality (Uosm), which has been commonly adopted to explain the physiology of osmotic responses, indicated that a near maximally concentrated urine ( $\sim 1100$  mmol/kg) should occur at a Posm of 292 mmol/kg and AVP value of 4.6 pg/mL (43). This result contrasts with everyday observations but is easily understood from a starting Uosm:Posm ratio of approximately

$$187:282 = 0.66 \quad (2)$$

which can be back calculated from a starting AVP value of 1 pg/mL in these experiments (43). If we assume unity between plasma and urine electrolyte concentrations and accept that urea contributes 40% to Uosm (77), any Uosm:Posm ratio  $\leq 1.5$  is consistent with electrolyte-free renal water clearance and a water-loaded state (77). In contrast, the change in Posm (12 mmol/kg) that is responsible for the 1100-mmol/kg Uosm is entirely consistent with a hyperosmolar state:

$$\begin{aligned} \Delta \text{Uosm} &= 250 \times 0.35 \Delta \text{Posm} \text{ or } \Delta 12 \text{ Posm} \\ &= \Delta 1050 \text{ Uosm (27)} \end{aligned} \quad (3)$$

Therefore, both aspects of osmoregulation (ie, the variation and basal set point) are very important considerations when Posm is used to assess dehydration. When a person's true Posm baseline is not known, biological variation analysis can provide confident probabilistic estimates of dehydration by using both single and serial measures of Posm.



**FIGURE 3.** Means [men (●); women (■)] from 4 basal plasma osmolality samples under conditions of ad libitum fluid intake in 15 healthy subjects. Boundaries of rectangular boxes represent osmotic thresholds for unequivocal thirst (top) and AVP secretion (bottom) determined from regression analysis during hypertonic saline infusion. Dashed lines represent group means often cited in the literature. Adapted from reference 13 with permission (copyright 1976; The Endocrine Society). AVP, arginine vasopressin;  $P_{\text{osm}}$ , plasma osmolality.




### Biological variation of Posm (single measure)

Claude Bernard's concept of the tightly regulated milieu intérieur is commonly referenced by extension as a narrow 1–2% variation in Posm. Surprisingly few studies have quantified the intraindividual variation in Posm from day to day, but they have been consistent with this concept and reported values that ranged from 0.8% to 1.4% (2–4 mmol/kg), with the exclusion of the analytic (measurement) variation ( $\sim 0.5\%$  or 1 mmol/kg) (54, 78, 79). These studies (54, 78, 79) were not stratified and, thus, included contributions to the variation because of sex (73) and age (74). Although most measures of physiologic interest have a larger interindividual variation than intraindividual variation (80), the 2 measures are similar for Posm. This outcome seems to contrast with wide population reference intervals until it is considered that the variation between subjects shrinks when measurement methodology and other preanalytic factors are controlled for (80). The ratio of intraindividual to interindividual variation (index of individuality) in Posm ranges from 0.9 to 1.4 (54, 78, 79). The index of individuality provides a statistical framework to distinguish pathologic states such as dehydration from a single measurement. Any atypical value for a given individual, relative to the larger population of individuals, will go unnoticed when the ratio is  $<0.6$  but will be captured when the ratio is  $>1.4$  (80, 81). The probability of identifying an atypical value increases rapidly as the ratio exceeds 0.60 and approaches unity (1.00) (81).

The index of individuality concept for Posm is shown in Figure 4 and includes Uosm and saliva osmolality (Sosm) for comparison (54, 58). The interindividual variation is depicted by the differences in means (dots), whereas the intraindividual differences (typical day-to-day variation) in body fluid measures are represented by the range (bars) that surrounds each individual mean. "X" values in Figure 4 represent body fluid measures in response to a  $-2.5\text{-L}$  loss of body water ( $-3\%$  dehydration). When graphed relative to the respective dehydration thresholds determined empirically by receiver operating characteristic curve analysis, probabilities of false-negative and -positive findings become apparent. The index of individuality for Posm was 0.90. For contrast, ratios for Uosm and Sosm were 0.49 and 0.27, respectively (54). As illustrated in Figure 4, an atypical value for Posm (X, dehydrated) is more easily and accurately detected than an atypical value for Uosm or Sosm, despite the expected linear associations commonly reported when the dehydration level against Sosm or Uosm is regressed. The complete biological variation analysis (54) supports a Posm threshold of  $301 \pm 5$  mmol/kg, which is mathematically identical to the  $-0.56^\circ\text{C}$  depression in the freezing point proposed by Olmstead et al (82)  $>50$  y ago as a positive test for hypernatremia. We recommend the inclusion of a variance term ( $\pm 5$  mmol/kg) to account for biological differences in basal set points (54) and note its consistency to values (295–300 mmol/kg) reached by consensus (24) as consistent with impending dehydration.

### Biological variation of Posm (serial measures)

Reference change values (80) allow the observation of serial changes in Posm to be interpreted in terms of diagnosing dehydration (54, 58). Reference change values can be calculated (when the proper statistical assumptions are met) (80, 83)




**FIGURE 4.** Means (●) and absolute ranges of plasma (A), urine (B), and saliva (C) osmolality from 3 samples taken from each of 18 normally hydrated (euhydrated) subjects. "X" values represent a single sample from the same subjects after dehydration to  $-3.3 \pm 0.6\%$  of body mass. Dehydration thresholds were determined by using receiver operating characteristic curve analysis (value that provided the highest diagnostic accuracy) with the use of both euhydrated and dehydrated matched pairs. Data are a composite from references 54 and 58. P<sub>osm</sub>, plasma osmolality; S<sub>osm</sub>, saliva osmolality; U<sub>osm</sub>, urine osmolality.

by using the sum of analytic and intraindividual variations in Posm (80). The probability that a measured change in Posm is atypical can then be determined (58, 80). As illustrated in Figure 5, atypical changes in Posm begin above the daily 2–4-mmol/kg constancy threshold and provide increasing predictive certainty that dehydration has occurred in accordance with the equation

$$\text{Probability} = 1 - e^{-0.327x} \quad (4)$$

where  $x$  is the measured change in Posm (58). As a result, the probability that a change in Posm reflects the occurrence of dehydration can be gauged in both quantitative and qualitative terms.



**FIGURE 5.** Changes in  $P_{\text{osm}}$  plotted as a function of the statistical probability for dehydration. Semantic descriptors provide a scale that aligned with statistical probability to communicate the likelihood of dehydration in simple terms. The equation (inset) describes the curved line and its ubiquitous  $K$  constant ( $-0.327$ ). From Cheuvront et al (58).  $P_{\text{osm}}$ , plasma osmolality.

### LIMITATIONS OF USE OF OSMOMETRY FOR ASSESSMENT OF DEHYDRATION

It is clear that  $P_{\text{osm}}$  is critical for body water regulation, and plasma is a unique body fluid for use in the assessment of dehydration. Although  $U_{\text{osm}}$  and  $S_{\text{osm}}$  have also been used successfully for this purpose (84, 85), human variation in these body fluids seem to limit their potential utility (Figure 4). The greater variation in  $U_{\text{osm}}$  and  $S_{\text{osm}}$  is not surprising. For example,  $\leq 40\%$  of  $U_{\text{osm}}$  is attributable to urea (compared with only  $\sim 1\%$  for  $P_{\text{osm}}$ ), and thus, the addition of solute in the form of antecedent diet or catabolic byproducts of protein metabolism associated with exercise or illness may increase  $U_{\text{osm}}$  by the addition of solute (86, 87). Similar limitations apply for urine-specific gravity, whereas all urine-concentration measures are subject to timing and uniformity concerns that manifest empirically as differences between first morning, 24-h, and spot urine measures (88) in addition to acute drinking and exercise behaviors (59). The discriminatory power of renal excretion measures for the detection of dehydration is clearly secondary to changes in  $P_{\text{osm}}$  (28), but this does not, in any way, minimize the critical use of  $U_{\text{osm}}$  (and its relation to  $P_{\text{osm}}$ ) in the measurement of renal function related to the phenomenologic interpretation or differential diagnosis of other disorders (5, 28, 29). With regard to saliva,  $S_{\text{osm}}$  is subject to practical use issues related to simple oral artifacts (54, 89).  $S_{\text{osm}}$  may also be affected by anything that affects salivation (salivary flow), which includes a multitude of factors (85). Limitations of the use of  $P_{\text{osm}}$  for the assessment of dehydration must also be acknowledged.

$P_{\text{osm}}$  and plasma tonicity (effective osmolality) are very similar quantities in health (90). However, substances in the blood that raise osmolality but not tonicity (ineffective or penetrating solutes) have the potential to confound dehydration assessment. The calculation of the osmol gap will reveal contributions from ineffective solutes, but the direct measurement of  $P_{\text{osm}}$  is always recommended for dehydration assessment because of the large acceptable error in calculated osmolality ( $\pm 10$  mmol/kg) (90). Fluctuations in the volume of body fluid compartments will also affect  $P_{\text{osm}}$ . For example, consumption of a large meal can increase  $P_{\text{osm}}$  because of the osmolar shift of water out of the vasculature and into the gut (91). In contrast, simple changes in posture (42, 92) and even low-intensity exercise ( $\leq 40\%$  maximal

oxygen uptake) (93, 94) produce little effect, probably because osmotic concordance is not disrupted by the 2-way fluid flow between interstitial and intravascular spaces that share the same osmotic pressures (61, 64, 93). Higher exercise intensities increase  $P_{\text{osm}}$  as a result of greater intravascular volume losses and the presence of lactic acid, but recovery appears complete in  $\leq 20$ –30 min (93). Finally, as stated earlier,  $P_{\text{osm}}$  is of no use for the detection of volume depletion. When this distinction is made, coupled with the importance of biological variation and other issues discussed herein, criticisms for adopting  $P_{\text{osm}}$  as a gold standard for dehydration assessment (15, 95–97) are minimal.

### CONCLUSIONS AND FUTURE DIRECTIONS

Dehydration is a common physiologic state with implications for health and performance (1–8). Although the physiology of dehydration is well described, it remains difficult to assess accurately in practice. In this review, we highlighted how the phenomenologic interpretation of dehydration depends critically on the type and magnitude of dehydration, which directly affect threshold osmotic and volume-dependent compensatory anti-diuretic and thirst responses. We also emphasized how knowledge of biological variation improves our broader understanding of the physiology that underpins the osmotic response to dehydration in humans and affords important diagnostic insight for dehydration assessment. To help improve the practice of dehydration assessment, a single, atypical  $P_{\text{osm}}$  threshold value of  $301 \pm 5$  mmol/kg is suggested (54) as a starting point for this purpose, along with a nomogram (58) for the estimation of the probability of dehydration when serial changes in  $P_{\text{osm}}$  are measured as an adjunct to quantitative differential diagnostic procedures. No standard method has been advocated for the assessment of volume depletion (16, 19), but a 20-beats/min sit-to-stand cutoff provides high test specificity for both dehydration and volume depletion (65, 69). Because  $P_{\text{osm}}$  requires the collection of blood and the preparation of plasma, future efforts to identify or develop an acceptable noninvasive surrogate for  $P_{\text{osm}}$  would benefit clinical, sports, and military medicine communities (7). A test with high diagnostic accuracy for moderate volume depletion is also needed.

We thank Brett R Ely, Mary C Pardee, and Kristen R Heavens for technical assistance with this manuscript.

The authors' responsibilities were as follows—all authors: conception, writing, and editing of the manuscript. None of the authors had a conflict of interest.

### REFERENCES

- Adrogue HJ, Madias NE. Hyponatremia. *N Engl J Med* 2000;342:1493–9.
- Campbell N. Dehydration: why is it still a problem? *Nurs Times* 2011; 107:12–5.
- Feig PU, McCurdy DK. The hypertonic state. *N Engl J Med* 1977;297:1444–54.
- King CK, Glass R, Bresee JS, Duggan C. Managing acute gastroenteritis among children: oral rehydration, maintenance, and nutritional therapy. *MMWR Recomm Rep* 2003;52:1–16.
- Nguyen MK, Kurtz I. A new quantitative approach to the treatment of the dysnatremias. *Clin Exp Nephrol* 2003;7:125–37.
- Rose BD. Hyperosmolar states-hyponatremia. In: Burton DR, ed. *Clinical physiology of acid-base and electrolyte disorders*. New York, NY: McGraw-Hill, 1994:695–736.

7. Institute of Medicine. Water. In: Dietary Reference Intakes. Washington, DC: National Academies Press, 2005:73–185.
8. Sawka MN, Burke LM, Eichner ER, Maughan RJ, Montain SJ, Stachenfeld NS. American College of Sports Medicine position stand. Exercise and fluid replacement. *Med Sci Sports Exerc* 2007;39:377–90.
9. Stookey JD, Pieper CF, Cohen HJ. Hypertonic hyperglycemia progresses to diabetes faster than normotonic hyperglycemia. *Eur J Epidemiol* 2004;19:935–44.
10. Stookey JD, Barclay D, Arief A, Popkin BM. The altered fluid distribution in obesity may reflect plasma hypertonicity. *Eur J Clin Nutr* 2007;61:190–9.
11. Black RE, Morris SS, Bryce J. Where and why are 10 million children dying every year? *Lancet* 2003;361:2226–34.
12. Warren JL, Bacon WE, Harris T, McBean AM, Foley DJ, Phillips C. The burden and outcomes associated with dehydration among US elderly, 1991. *Am J Public Health* 1994;84:1265–9.
13. Robertson GL. The regulation of vasopressin function in health and disease. *Recent Prog Horm Res* 1976;33:333–85.
14. Bourque CW. Central mechanisms of osmosensation and systemic osmoregulation. *Natl Rev Neurosci* 2008;9:519–31.
15. Armstrong LE. Assessing hydration status: the elusive gold standard. *J Am Coll Nutr* 2007;26(5 suppl):575S–84S.
16. Peacock WF, Soto KM. Current techniques of fluid status assessment. *Contrib Nephrol* 2010;164:128–42.
17. Crecelius C. Dehydration: myth and reality. *J Am Med Dir Assoc* 2008;9:287–8.
18. Hayajneh WA, Jdaitawi H, Al SA, Hayajneh YA. Comparison of clinical associations and laboratory abnormalities in children with moderate and severe dehydration. *J Pediatr Gastroenterol Nutr* 2010;50:290–4.
19. McGee S, Abernethy WB III, Simel DL. The rational clinical examination. Is this patient hypovolemic? *JAMA* 1999;281:1022–9.
20. Robinson BE, Weber H. Dehydration despite drinking: beyond the BUN/Creatinine ratio. *J Am Med Dir Assoc* 2004;5:S67–71.
21. Roland D, Clarke C, Borland ML, Pascoe EM. Does a standardised scoring system of clinical signs reduce variability between doctors' assessments of the potentially dehydrated child? *J Paediatr Child Health* 2010;46:103–7.
22. Steiner MJ, DeWalt DA, Byerley JS. Is this child dehydrated? *JAMA* 2004;291:2746–54.
23. Steiner MJ, Nager AL, Wang VJ. Urine specific gravity and other urinary indices: inaccurate tests for dehydration. *Pediatr Emerg Care* 2007;23:298–303.
24. Thomas DR, Cote TR, Lawhorne L, Levenson SA, Rubenstein LZ, Smith DA, Stefanacci RG, Tangelos EG, Morley JE. Understanding clinical dehydration and its treatment. *J Am Med Dir Assoc* 2008;9:292–301.
25. Vivanti A, Harvey K, Ash S, Battistutta D. Clinical assessment of dehydration in older people admitted to hospital: what are the strongest indicators? *Arch Gerontol Geriatr* 2008;47:340–55.
26. Reeves WB, Bichet DG, Andreoli TE. Posterior pituitary and water metabolism. In: Wilson JD, Foster DW, Kronenberg HM, Larsen PR, eds. *Williams textbook of endocrinology*. 9th ed. Philadelphia, PA: WB Saunders Co, 1998:341–87.
27. Robertson GL, Shelton RL, Athar S. The osmoregulation of vasopressin. *Kidney Int* 1976;10:25–37.
28. Shoker AS. Application of the clearance concept to hyponatremic and hypernatremic disorders: a phenomenological analysis. *Clin Chem* 1994;40:1220–7.
29. Haraway AW Jr, Becker EL. Clinical application of cryoscopy. *JAMA* 1968;205:506–12.
30. Mange K, Matsuura D, Cizman B, Soto H, Ziyadeh FN, Goldfarb S, Neilson EG. Language guiding therapy: the case of dehydration versus volume depletion. *Ann Intern Med* 1997;127:848–53.
31. Armstrong LE. Hydration assessment techniques. *Nutr Rev* 2005;63:S40–54.
32. Oppliger RA, Bartok C. Hydration testing of athletes. *Sports Med* 2002;32:959–71.
33. Antunes-Rodrigues J, de Castro M, Elias LLK, Valença MM, McCann SM. Neuroendocrine control of body fluid metabolism. *Physiol Rev* 2004;84:169–208.
34. Gonzalez RR, Chevront SN, Montain SJ, Goodman DA, Blanchard LA, Berglund LG, Sawka MN. Expanded prediction equations of human sweat loss and water needs. *J Appl Physiol* 2009;107:379–88.
35. Sawka MN, Chevront SN, Carter R III. Human water needs. *Nutr Rev* 2005;63:S30–9.
36. de Castro JM. A microregulatory analysis of spontaneous fluid intake by humans: evidence that the amount of liquid ingested and its timing is mainly governed by feeding. *Physiol Behav* 1988;43:705–14.
37. Shore AC, Markandu ND, Sagnella GA, Singer DR, Forsling ML, Buckley MG, Sugden AL, MacGregor GA. Endocrine and renal response to water loading and water restriction in normal man. *Clin Sci (Lond)* 1988;75:171–7.
38. Phillips PA, Rolls BJ, Ledingham JG, Morton JJ. Body fluid changes, thirst and drinking in man during free access to water. *Physiol Behav* 1984;33:357–63.
39. Segar WE, Moore WW. The regulation of antidiuretic hormone release in man. *J Clin Invest* 1968;47:2143–51.
40. Dunn FL, Brennan TJ, Nelson AE, Robertson GL. The role of blood osmolality and volume in regulating vasopressin secretion in the rat. *J Clin Invest* 1973;52:3212–9.
41. Robertson GL, Athar S. The interaction of blood osmolality and blood volume in regulating plasma vasopressin in man. *J Clin Endocrinol Metab* 1976;42:613–20.
42. Kimura T, Minai K, Matsui K, Mouri T, Sato T. Effect of various states of hydration on plasma ADH and renin in man. *J Clin Endocrinol Metab* 1976;42:79–87.
43. Robertson GL, Mahr EA, Athar S, Sinha T. Development and clinical application of a new method for the radioimmunoassay of arginine vasopressin in human plasma. *J Clin Invest* 1973;52:2340–52.
44. Nadai JW, Pedersen S, Maddock WG. A comparison between dehydration from salt loss and from water deprivation. *J Clin Invest* 1941;20:691–703.
45. Baylis PH, Robertson GL. Plasma vasopressin response to hypertonic saline infusion to assess posterior pituitary function. *J R Soc Med* 1980;73:255–60.
46. Zerbe RL, Miller JZ, Robertson GL. The reproducibility and heritability of individual differences in osmoregulatory function in normal human subjects. *J Lab Clin Med* 1991;117:51–9.
47. Greenleaf JE, Morimoto T. Mechanisms controlling fluid ingestion: thirst and drinking. In: Buskirk ER, Puhl SM, eds. *Body fluid balance: exercise and sport*. Boca Raton, FL: CRC Press, 1996:3–17.
48. Geelen G, Keil LC, Kravik SE, Wade CE, Thrasher TN, Barnes PR, Pyka G, Nesvig C, Greenleaf JE. Inhibition of plasma vasopressin after drinking in dehydrated humans. *Am J Physiol* 1984;247:R968–71.
49. Denton D, Shade R, Zamarippa F, Egan G, Blair-West J, McKinley M, Fox P. Correlation of regional cerebral blood flow and change of plasma sodium concentration during genesis and satiation of thirst. *Proc Natl Acad Sci USA* 1999;96:2532–7.
50. Greenleaf JE, Sargent F. Voluntary dehydration in man. *J Appl Physiol* 1965;20:719–24.
51. Rothstein A, Adolph EF, Wills JH. Voluntary dehydration. In: Visscher MB, Bronk DW, Landis EM, Ivy AC, eds. *Physiology of man in the desert*. Rochester, NY: Interscience Publishers Inc, 1947:254–77.
52. Gudivaka R, Schoeller DA, Kushner RF, Bolt MJ. Single- and multi-frequency models for bioelectrical impedance analysis of body water compartments. *J Appl Physiol* 1999;87:1087–96.
53. Chevront SN, Carter R III, Montain SJ, Sawka MN. Daily body mass variability and stability in active men undergoing exercise-heat stress. *Int J Sport Nutr Exerc Metab* 2004;14:532–40.
54. Chevront SN, Ely BR, Kenefick RW, Sawka MN. Biological variation and diagnostic accuracy of dehydration assessment markers. *Am J Clin Nutr* 2010;92:565–73.
55. Greenleaf JE. Problem: thirst, drinking behavior, and involuntary dehydration. *Med Sci Sports Exerc* 1992;24:645–56.
56. Johnson RE. Water and osmotic economy on survival rations. *J Am Diet Assoc* 1964;45:124–9.
57. Bartok C, Schoeller DA, Sullivan JC, Clark RR, Landry GL. Hydration testing in collegiate wrestlers undergoing hypertonic dehydration. *Med Sci Sports Exerc* 2004;36:510–7.
58. Chevront SN, Fraser CG, Kenefick RW, Ely BR, Sawka MN. Reference change values for monitoring dehydration. *Clin Chem Lab Med* 2011;49:1033–7.
59. Popowski LA, Oppliger RA, Patrick LG, Johnson RF, Kim JA, Gisolf CV. Blood and urinary measures of hydration status during progressive acute dehydration. *Med Sci Sports Exerc* 2001;33:747–53.



60. Carter R III, Cheuvront SN, Vernieuw CR, Sawka MN. Hypohydration and prior heat stress exacerbates decreases in cerebral blood flow velocity during standing. *J Appl Physiol* 2006;101:1744–50.
61. Nose H, Mack GW, Shi XR, Nadel ER. Shift in body fluid compartments after dehydration in humans. *J Appl Physiol* 1988;65:318–24.
62. Sawka MN, Toner MM, Francesconi RP, Pandolf KB. Hypohydration and exercise: effects of heat acclimation, gender, and environment. *J Appl Physiol* 1983;55:1147–53.
63. Cheuvront SN, Kenefick RW, Montain SJ, Sawka MN. Mechanisms of aerobic performance impairment with heat stress and dehydration. *J Appl Physiol* 2010;109:1989–95.
64. Coleman TG, Norman RA, Manning RD. Equilibrium between extracellular and intracellular fluid. In: Guyton AC, Taylor AE, Granger HJ, eds. *Circulatory physiology II: dynamics and control of the body fluids*. Philadelphia, PA: WB Saunders Company, 1975:225–42.
65. Cheuvront SN, Ely BR, Kenefick RW, Buller MJ, Charkoudian N, Sawka MN. Hydration assessment using the cardiovascular response to standing. *Eur J Appl Physiol* 2012;112:4081–9.
66. Hoyt RW, Honig A. Body fluid and energy metabolism at high altitude. In: Fregley MJ, Blatteis CM, eds. *Handbook of physiology*. Vol 2, Sect 4. New York, NY: Oxford University Press, 1996:1277–89.
67. Lennquist S. Cold-induced diuresis. A study with special reference to electrolyte excretion, osmolal balance and hormonal changes. *Scand J Urol Nephrol* 1972;9(suppl 9):1–142.
68. O'Brien C, Young AJ, Sawka MN. Hypohydration and thermoregulation in cold air. *J Appl Physiol* 1998;84:185–9.
69. Witting MD, Gallagher K. Unique cutpoints for sitting-to standing orthostatic vital signs. *Am J Emerg Med* 2003;21:45–7.
70. Knopp R, Claypool R, Leonardi D. Use of the tilt test in measuring acute blood loss. *Ann Emerg Med* 1980;9:72–5.
71. Cooke WH, Ryan KL, Convertino VA. Lower body negative pressure as a model to study progression to acute hemorrhagic shock in humans. *J Appl Physiol* 2004;96:1249–61.
72. Fortes MB, Diment BC, Di FU, Gunn AE, Kendall JL, Esmaelpour M, Walsh NP. Tear fluid osmolality as a potential marker of hydration status. *Med Sci Sports Exerc* 2011;43:1590–7.
73. Stachenfeld NS, Splenser AE, Calzone WL, Taylor MP, Keefe DL. Sex differences in osmotic regulation of AVP and renal sodium handling. *J Appl Physiol* 2001;91:1893–901.
74. Kenney WL, Chiu P. Influence of age on thirst and fluid intake. *Med Sci Sports Exerc* 2001;33:1524–32.
75. Sollanek KJ, Kenefick RW, Cheuvront SN, Axtell RS. Potential impact of a 500-mL water bolus and body mass on plasma osmolality dilution. *Eur J Appl Physiol* 2011;111:1999–2004.
76. Zerbe RL, Robertson GL. Osmoregulation of thirst and vasopressin secretion in human subjects: effect of various solutes. *Am J Physiol* 1983;244:E607–14.
77. Shimizu K, Kurosawa T, Sanjo T, Hoshino M, Nonaka T. Solute-free versus electrolyte-free water clearance in the analysis of osmoregulation. *Nephron* 2002;91:51–7.
78. Fraser CG, Cummings ST, Wilkinson SP, Neville RG, Knox JD, Ho O, MacWalter RS. Biological variability of 26 clinical chemistry analytes in elderly people. *Clin Chem* 1989;35:783–6.
79. Jahan A, Browning MC, Fraser CG. Desirable performance standards for assays of serum water and osmolality. *Clin Chem* 1988;34:995.
80. Fraser CG. *Biological variation: from principles to practice*. Washington, DC: American Association of Clinical Chemistry Press, 2001.
81. Harris EK. Effects of intra- and interindividual variation on the appropriate use of normal ranges. *Clin Chem* 1974;20:1535–42.
82. Olmstead EG, Roth DA. The use of serum freezing point depressions in evaluating salt and water balance in preoperative and postoperative states. *Surg Gynecol Obstet* 1958;106:41–8.
83. Harris EK, Brown SS. Temporal changes in the concentrations of serum constituents in healthy men. Distributions of within-person variances and their relevance to the interpretation of differences between successive measurements. *Ann Clin Biochem* 1979;16:169–76.
84. Shirreffs SM, Maughan RJ. Urine osmolality and conductivity as indices of hydration status in athletes in the heat. *Med Sci Sports Exerc* 1998;30:1598–602.
85. Walsh NP, Montague JC, Callow N, Rowlands AV. Saliva flow rate, total protein concentration and osmolality as potential markers of whole body hydration status during progressive acute dehydration in humans. *Arch Oral Biol* 2004;49:149–54.
86. Meroney WH, Rubini ME, Blythe WB. The effect of antecedent diet on urine concentrating ability. *Ann Intern Med* 1958;48:562–73.
87. Pepper OH. Studies on the specific gravity of the urine. *J Clin Invest* 1924;1:13–9.
88. Gowans EM, Fraser CG. Despite correlation, random spot and 24-h urine specimens are not interchangeable. *Clin Chem* 1987;33:1080–1.
89. Ely BR, Cheuvront SN, Kenefick RW, Sawka MN. Limitations of salivary osmolality as a marker of hydration status. *Med Sci Sports Exerc* 2011;43:1080–4.
90. Kruse JA, Cadnapaphornchai P. The serum osmole gap. *J Crit Care* 1994;9:185–97.
91. Gill GV, Baylis PH, Flear CT, Lawson JY. Changes in plasma solutes after food. *J R Soc Med* 1985;78:1009–13.
92. Shirreffs SM, Maughan RJ. The effect of posture change on blood volume, serum potassium and whole body electrical impedance. *Eur J Appl Physiol Occup Physiol* 1994;69:461–3.
93. Greenleaf JE, Van BW, Brock PJ, Morse JT, Mangseth GR. Plasma volume and electrolyte shifts with heavy exercise in sitting and supine positions. *Am J Physiol* 1979;236:R206–14.
94. Montain SJ, Laird JE, Latzka WA, Sawka MN. Aldosterone and vasopressin responses in the heat: hydration level and exercise intensity effects. *Med Sci Sports Exerc* 1997;29:661–8.
95. Francesconi RP, Hubbard RW, Szlyk PC, Schnakenberg D, Carlson D, Leva N, Sils I, Hubbard L, Pease V, Young J, et al. Urinary and hematologic indexes of hypohydration. *J Appl Physiol* 1987;62:1271–6.
96. Hackney AC, Coyne JT, Pozos R, Feith S, Seale J. Validity of urine-blood hydration measures to assess total body water changes during mountaineering in the sub-Arctic. *Arctic Med Res* 1995;54:69–77.
97. Perrier E, Vergne S, Klein A, Poupin M, Rondeau P, Le Bellego L, Armstrong LE, Lang F, Stookey J, Tack I. Hydration biomarkers in free-living adults with different levels of habitual fluid consumption. *Br J Nutr* 2012;Aug 31 (Epub ahead of print).